

KENYAN COMMUNITY HEALTH PROFILE 2022



A BOLDER HEALTHIER BIRMINGHAM






WORKING AND LEARNING WELL

ACADEMIC ATTAI

Using the black and Indian ethnic categories as proxy to gain an insight into the Kenvan community on education attainment reveals, that black African pupils achieved Progress 8 scores higher than average (0.17). Indian pupils achieved the SECOND HIGHEST PROGRESS 8 SCORE (0.71)

ECONOMIC INACTIVITY *****18.9% *****30.5%



resident in the West Midlands own a property; ether the property is owned outright or with a mortgage/ loan or shared ownership.

UNEMPI NYMFN *****5.9% *****4.4%

A BOLDER HEALTHIER BIRMINGHAM

PROTECT AND DETECT

CANCER SCREENING

There is no data or information on the Kenyan community's take up of screening in the UK. However, research on uptake of Pap smear testing among Kenyan migrants in the Netherlands found willingness to participate was hindered by barriers such as

LANGUAGE. **PREFERENCE FOR FEMALE GPs.** ACCULTURATION. **VIEWS ABOUT THE HEALTH CARE SYSTEM**

SEXUAL HEAI TH

Research has found the communities in Britain most affected by the AIDS/HIV pandemic were nationals of

KENYA, UGANDA, ZAMBIA, ZIMBABWE, AND CONGO

TUBERCULOSIS (TB) IN 2020, PEOPLE BORN IN KENYA ACCOUNTED FOR 46 CASES

> of the UK's TB cases with a median time of 17 years from entry to notification since arrival to the UK



CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD) There is an absence of statistics on the prevalence of COPD

within the Kenyan community in the UK. In Kenya, chronic respiratory diseases are responsible for approximately

THE MAIN DRIVERS OF COPD INCLUDE **TOBACCO SMOKING. INDOOR AIR** POLLUTION. OUTDOOR AIR POLLUTION. **OCCUPATIONAL DUST AND CHEMICALS**

1.92% 1.5%

AGEING AND DYING WELL

Published research from the United States with Kenyan participants (n=51) found that among six African immigrant groups. Kenvans were more likely to report having diabetes

CARDIOVASCULAR DISEASE

The leading CVD deaths are stroke (6.1%.

diseases (4.6%, male 47%, female 4.6%)

male 5.8%, female 6.4%) and Ischemic Heart

IT IS ESTIMATED THAT MORTALITY

DUE TO CVD IN KENYA IS

.8% END

Research which included Kenvan participants (8%) in the study sample found that people with dementia and

their carers usually begin help-seeking from close family and then follow this up by consulting primary care physicians

DEMENTIA

Black African and Caribbean elders (BACE) have a higher prevalence and earlier onset of dementia compared with the indigenous white UK population

CLOSING THE GAPS

DEPRIVATION O AS ALMOST C

of Kenvans in the West Midlands either fully own a property or have a mortgage, and also have very low levels of unemployment, it is unlikely the Kenyan community in the UK faces deprivation